# **TEBIZOLE-25% EC**





Systemic fungicide with protective, curative action to control many fungal diseases on a wide range of crops.

**COMPOSITION:** Each liter contains:

TEBUCONAZOLE 25% W/V (A.I)

PROPERTIES:

**TEBIZOLE-25% EC:** Systemic fungicide with protective and curative action.

TEBIZOLE-25% EC: Fungicide recommended to control disease such as powdery mildew,

yellow rust, brown rust, septoria, scab, barley leaf spot.

#### USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE	NUMBER OF TREATMENS PER SEASON	INTERVAL BETWEEN TREATMENS	TREATMENTPERIOD	
					DAYS	GROWTH STAGE
Wheat	Powdery Mildew, yellow Rust, brown Rust, septorian leaf spots.	0.2-1.0 lt./ha	2	2 weeks	35	69
Barley	Powdery Mildew, scab, Leafspot	0.2-1.0 lt./ha	2	2 weeks	42	65
Oats	Powderymildew, rusts , Leaf spot.	0.2-1.0 lt./ha	2	2 weeks	42	65
Rye	Rusts	0.2-1.0 lt./ha	2	2 weeks	35	69

### CAUTION:

- · Used only on above mentioned crops and follow the recommended safety period before harvest.
- · Lowest dosages are used for early application of fungal development.

## **RE-ENTRY PERIOD:**

48 hours from last application.

#### COMPATIBILITY:

Compatible with most common used pesticides and compatibility test is recommended before application.

<sup>•</sup> For more details about first aid & precautions please refer to first aid & precautions index.