BIFENAMIN-10% EC

INSECTICIDE / ACARICIDE EMULSIFIABLE CONCENTRATE



Highly effective insecticide and Acaricide with contact and stomach action. It can be used as a protective treatment when applied at regular intervals or a knockdown treatment to control existing pests.

COMPOSITION: Each liter contains:

BIFENTHRIN 10% W/V (A.I)

PROPERTIES:

BIFENAMIN-10% EC: Is a pyrethroidal Acaricide /Insecticide which acts on the nervous system

of insects.

BIFENAMIN-10% EC: Effective against a broad range of insects on the agricultural crops in the

table of uses.

USES AND APPLICATION RATE:

CROPS	INSECT PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)	COMMENTS
Faba Beans, Clover, Barley, Field Peas, Lupine, Wheat	Mites, Cutworm.	1-2	4 Weeks (grazing)	
Peaches, Nectarines, Plums, Apricots	Beetles	10	1	There must be a minimum of 10 days between the re-treatment and the initial application.
Citrus	Weevil Larvae	Pre-emergence program 12.5 or 25 ml/tree		Apply a total spray volume of 5-10 lt. / tree Pre-emergence program: apply just prior to or at the first sign of major beetle emergence in mid-october.
		Post-emergence Monitoring program 6 ml/tree		Post-emergence program: apply at beetle emergence in october/ november. Follow up treatment maybe necessary based on a threshold of 25 beetles per 10 sites per orchard.

[•] For more details about first aid & precautions please refer to first aid & precautions index.

CROPS	INSECT PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)	COMMENTS
Grapes	Beetle larve	200		The application must be made at late dormancy after pruning and before budburst.
Pears	Mealybug	5 ml with SAMAROL (summer oil) at (1L /100L)	14	Do not spray until young nymphs emerge in spring.
Tomatoes	Mites, Worms	12	1	Do not apply this product to helicoverpa armigera larvae larger than 5 mm in length.
	Whitefly	6		

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours from last application.

COMPATIBILITY:

Compatible with most pesticides except alkaline compounds, but compatibility test is recommended before spraying.

[•] For more details about first aid & precautions please refer to first aid & precautions index.