Mn-PROVAP

FERTILIZER SOLUBLE LIQUID



COMPOSITION: Each It. Contains:

Manganese 3.0% W/V (A.I)

PROPERTIES:

MN-PROVAP: used for the prevention and correction of manganese deficiency in crops listed on

the table of uses.

MN-PROVAP: it is used as a foliar spray.

USES AND APPLICATION RATE:

CROPS	APPLICATION RATE (ML\DONUM)	APPLICATION TIMING	APPLICATION INTERVAL (DAYS)
Potato	118.29 -946.35	At tuber initiation	14-21
Onion	118.29 - 473.16	After thinning, or 3- to 4-leaf stage	
Celery, Lettuce	118.29 - 473.16		
Broccoli, Cabbage, Cauliflower	118.29 - 473.16		
Beans, Soybeans	118.29 -473.16	From the appearance of the trifoliate leaf at the 5 node until its appearance at the 6 node and branching	14-21 until senescence
Peppers, Tomato, Eggplant	118.29 - 473.16	At crop establishment	14-21
Melons, Cucumber, Squash	118.29 -946.35	At 3- to 4-leaf stage	14-21 until harvest
Oranges, Lemons	236.52 -709.74	At first bloom	
Apple, Pear	118.29 -946.35		
Cherry, Peach, Plum	118.29 -946.35		
Strawberry, Blueberry	118.29 -946.35		
Grape	118.29 - 473.16	New shoot growth height 10.16-15.24 cm	
Almond, Hazelnut, Pistachio, Pecan, Walnut	236.52 -709.74	At first bloom	14-21 until harvest
Cereal	118.29 - 473.16	Begin on emergence of spring growth or deficiency symptoms.	14-21
Grain, Forage, Fodder, and Straw	118.29 - 473.16		

[·] For more details about first aid & precautions please refer to first aid & precautions index.

CROPS	APPLICATION RATE (ML\DUNUM)	APPLICATION TIMING	APPLICATION INTERVAL (DAYS)
Animal Feed, Non Grass	118.29 -946.35	At 2- to 4-leaf stage	
Cotton, Sunflower	118.29 - 473.16	A 2- to 4-leaf stage	14-21
Avocado, Artichoke, Hops, Flax, Mint, Olive, Peanut, Tobacco	118.29 -946.35	At first bloom	14-21 until harvest

CAUTION:

- Used only the recommended application rate

PHYTOTOXICITY:

Not phytotoxic when use as recommended. Treat a small group of target crop including plant cultivars and evaluate for 3 to 7 days prior to making an application to your entire crop.

COMPATIBILITY:

Compatible with most pesticides and fertilizers, when mixed with phosphate fertilizers vigorous agitation is required and compatibility test is recommended before application and it's preferable to be used after preparation.

[·] For more details about first aid & precautions please refer to first aid & precautions index.