



NEXIFLOR
 (Injectable Solution)
 For cattle

COMPOSITION: Each 1 ml contains:

| | |
|---|----------------|
| Florfenicol | 300 mg |
| Flunixin (as flunixin meglumine) | 16.5 mg |

INDICATION:

Treatment of respiratory infections caused by Mannheimia haemolytica, Pasteurella multocida, Mycoplasma bovis and Histophilus somni associated with pyrexia.

DOSAGE AND ADMINISTRATION:

- Single subcutaneous injection.
- 2 mL/15 kg body weight.
- ✓ The dose volume given at any one injection site should not exceed 10 ml.
- ✓ It is recommended to treat animals in the early stages of the disease and to evaluate the response to treatment 48 hours after injection.
- ✓ The anti-inflammatory component, flunixin, may mask a poor bacteriological response to florfenicol in the first 24 hours after injection. If clinical signs of respiratory disease persist or increase, or if relapse occurs, treatment should be changed, using another antibiotic, and continued until clinical signs have resolved.

OVERDOSE

- The overdose in animals may cause decrease in food and water consumption which affect the weight gain of the animals.
- Erosive and ulcerative abomasum lesions.



VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



CONTRA-INDICATIONS

- Do not use in adult bulls intended for breeding purposes.
- Do not use in animals suffering from hepatic, cardiac and renal diseases.
- Do not use if there is a risk of gastrointestinal bleeding or in cases where there is evidence of altered hemostasis.
- Do not use in the case of known hypersensitivity to active.

ADVERSE REACTIONS

In the injection site, may result a swelling that becomes palpable 2-3 days after injection.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

The product must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Gastrointestinal tract ulceration may be exacerbated by corticosteroids in animals given NSAIDs

WITHDRAWAL PERIOD:

Meat and offal: 46 days.

Milk: Not authorized for use in lactating animals producing milk for human consumption.

Do not use during lactation or drying off periods. Do not use in pregnant animals which are intended to produce milk for human consumption within 2 months of expected parturition

INCOMPATIBILITIES

Do not mix the product with other medicinal products.

STORAGE:

Store at a temperature below 30° C.

Keep out of reach and sight of children.

For VET. Use Only