



CLOMEVAP

(Suspension)

For Sheep and lambs.

COMPOSITION: Each ml contains:

Closantel	50 mg
Mebendazole	75 mg

INDICATIONS:

For the treatment and control of liver fluke, gastro-intestinal nematodes, lungworms, cestodes and larval stages of some arthropods.

Trematodes:

Liver flukes:

Fasciola hepatica (adults + 5-8 week immatures)

Fasciola gigantica (adults + 8 week immatures)

Nematodes:

Roundworms:

Haemonchus contortus, Bunostomum spp., Chabertia ovina, Gaigeria pachyscelis, Oesophagostomum spp., Capillaria spp., Cooperia spp., Nematodirus spp., Ostertagia circumcincta, Trichostrongylus axei, Trichostrongylus colubriformis, Trichostrongylus vitrines, Trichuris ovis, Strongyloides papillosus.

Lungworms:

Dictyocaulus filarial.

Cestodes:

Avitellina spp., Moniezia spp.

Arthropods:

Oestrus ovis (nasal bot), Ticks (Ixodes ricinus) feeding on sheep at the time of treatment are likely to produce fewer viable eggs.



DOSAGE AND ADMINISTRATION ROUTE:

- For oral administration by drenching gun.
- Shake well before use.
- The recommended dose is 1 ml per 5 kg BW.
- All sheep on infested pasture should be dosed at regular intervals during the fluke season. The interval between dosing will depend on the level of pasture contamination; in severe fluke seasons, dosing every 6-8 weeks may be necessary. Is active against worm eggs and prevents pasture contamination with fluke eggs for approximately 13 weeks.
- Treatment intervals of 10-12 weeks throughout the fluke season are recommended.

OVERDOSE:

Symptoms of acute closantel overdosage are decreased vision or blindness, anorexia, incoordination and general weakness.

WITHDRAWAL PERIOD

Meat and offal: 65 days.

Not authorized for use in ewes producing milk for human consumption including during the dry period. Do not use within 1 year prior to the first lambing in ewes intended to produce milk for human consumption.

STORAGE: Store in dry place below 30 °C.

For vet use only