



# CLOMEVAP

(Oral Suspension)  
For Sheep and lambs.

## COMPOSITION: Each ml contains:

<b>Closantel</b>	<b>50 mg</b>
<b>Mebendazole</b>	<b>75 mg</b>

## INDICATIONS:

For the treatment and control of liver fluke, gastro-intestinal nematodes, lungworms, cestodes and larval stages of some arthropods.

- **Trematodes:**

- **Liver flukes:**

- Fasciola hepatica (adults + 5–8-week immatures)

- Fasciola gigantica (adults + 8-week immatures)

- **Nematodes:**

- **Roundworms:** Haemonchus contortus, Bunostomum spp., Chabertia ovina, Gaigeria pachyscelis, Oesophagostomum spp., Capillaria spp., Cooperia spp., Nematodirus spp., Ostertagia circumcincta, Trichostrongylus axei, Trichostrongylus colubriformis, Trichostrongylus vitrines, Trichuris ovis, Strongyloides papillosus.

- **Lungworms:** Dictyocaulus filarial.

- **Cestodes:** Avitellina spp., Moniezia spp.

- **Arthropods:** Oestrus ovis (nasal bot), Ticks (Ixodes ricinus) feeding on sheep at the time of treatment are likely to produce fewer viable eggs.



## **DOSAGE AND ADMINISTRATION ROUTE:**

- For oral administration by drenching gun.
- Shake well before use.
- The recommended dose is 1 ml per 5 kg BW.
- All sheep on infested pasture should be dosed at regular intervals during the fluke season. The interval between dosing will depend on the level of pasture contamination; in severe fluke seasons, dosing every 6-8 weeks may be necessary. Clomevap Is active against worm eggs and prevents pasture contamination with fluke eggs for approximately 13 weeks.
- Treatment intervals of 10-12 weeks throughout the fluke season are recommended.

## **OVERDOSE:**

Symptoms of acute closantel overdosage are decreased vision or blindness, anorexia, incoordination and general weakness.

## **WITHDRAWAL PERIOD**

**Meat and offal:** 65 days.

- Not authorized for use in ewes producing milk for human consumption including during the dry period. Do not use within 1 year prior to the first lambing in ewes intended to produce milk for human consumption.

**STORAGE:** Store in dry place below 30 °C.

**PACKING:** (100, 250, 500, 1000) ml.

**FOR VET USE ONLY**