



# VAPCO

## VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.



# FIPROVAP – 1%

Pour-On  
Tick killer and Insecticide

**COMPOSITION: Each ml contains:**

**Fipronil** **10 mg**

### DESCRIPTION:

Treatment and control of horn fly, ticks, Dermatobia (human bot fly), chewing and sucking louse, and myiasis.

Is an insecticide which active ingredient is fipronil, a member of a new class of chemicals known as phenylpyrazole that controls external parasites resistant to pyrethroids, organophosphates and amidines. Due to its formulation (low volume and ready to use) when is applied on animal's dorsal line, it gets a distribution all over the skin and hair of animals because it has an affinity for sebaceous glands and its secretions.

### BENEFITS:

High efficacy against ticks and horn fly strains resistant to pyrethroids and organophosphates.  
Easy application. Major control for longer time.  
Less stress due to less movements and baths. More weight gain. Safety for animal and person who handles it.

### INDICATIONS:

Treatment and control of the following external parasites of **cattle, sheep, camelids, horses, guinea pigs**: Ticks (*Boophilus microplus*), horn fly (*Haematobia irritans*), chewing lice (*Damalinea bovis*), sucking lice (*Linognathus vituli*) and nodular myiasis (*Dermatobia hominis*); flies (*Melophagus ovinus*) and **poultry**: fleas (*Echidnophaga gallinacea*), louses (*Menopon gallinae*, *Lipeurus caponis*) and mites (principally *Dermanyssus gallinae* or red mite).

### DOSAGE AND APPLICATION METHOD:

Epicutaneous application (pour-on).

- ✓ **Cattle, sheep, camelids and horses:** The recommended dose is 1 mL/10 kg of body weight, equivalent to 1 mg of fipronil/kg b.w.
- ✓ **Guinea pigs and poultry:** The recommended dose is of 0.1 mL/kg b.w., equivalent to 1 mg of fipronil/kg b.w.; which in the practice is equivalent to two drops per each kg b.w., applied on the back of the animal.
- Apply along the dorsal line in a narrow strip extending along the top of the back from the withers to the tailhead.



- For optimal results, **Fiprovap** must be applied to all the animals in the herd on an external parasitic control program.
- The treatment interval could be at least every two months, it will depend on the severity of the infestation, weather and management conditions in the herd. Consult your veterinarian to determine the better moment for the treatment.

### WITHDRAWAL PERIOD:

Meat: 100 days.

Do not use in lactating cattle producing milk for human consumption.

Do not use in pregnant cows or heifers, which are intended to produce milk for human consumption, within 3 months of expected parturition.

### USE RESTRICTIONS:

External use only. Do not apply by another route of administration.

### CONTRAINDICATIONS, NON-DESIRED EFFECTS AND PRECAUTIONS:

- It does not present secondary effects or incompatibilities.
- Apply the product to the animal in a ventilated area.
- Do not use in sick or convalescent animals.
- Keep in a dry, cool place, protected from light. Storage below 30° C.
- Do not apply the product to wounded skin.

### USE PRECAUTIONS:

- Do not reuse the empty container.
- Avoid contaminating leaks such as rivers, streams, etc.
- Wash the exposed hands and skin after its use. If there is a direct contact with the skin, wash with plenty of water and soap.
- If there is a contact with the eyes, do it with plenty of water for 15 minutes.
- If ingested, drink large amounts of water, going immediately to the doctor with this leaflet.
- If there is a case of intoxication, take the person immediately to emergency and show this leaflet to the doctor.
- There is not a specific antidote.

**STORAGE:** Keep in dry, dark place below 30°C.

**PACKING:** (100, 250, 500, 1000) ml.

## FOR VET USE ONLY