



# VAPCO

**VETERINARY & AGRICULTURAL  
PRODUCTS MANUFACTURING CO. LTD.**



## NEOVAP

**Solution**

**For Cattle, Sheep, Goats and Poultry**

**COMPOSITION: Each ml contains:**

**Neomycin Sulphate                      200 mg**  
**(Equivalent to neomycin base 140 mg)**

**INDICATIONS:**

Neovap is indicated for the treatment diarrhea and control of colibacillosis caused by E. coli susceptible to neomycin in Cattle, Sheep, Goat and Poultry and bacterial enteritis.

**DOSAGE:**

Treatment of colibacillosis should continue for 14 days as follow:

<b>Kg (Body weighy)</b>	<b>Volume of Neovap</b>
Up to 10 kg	1 ml
Up to 50 kg	2.5 ml
Up to 100 kg	5 ml
Up to 300 kg	15 ml
More than 300 kg	30 ml

**Poultry dose:**

- From 1 day to 6 weeks: 50-100 ml/200 liter of water.
- More than 6 weeks: 100 ml/200 liter of water.
- Treatment for 3-5 days.
- ✓ In divided dose in drinking water to be consumed in 12-24 hours.
- ✓ It may be administered daily as drench for animals that are not drinking or eating.

ISO 9001 & 14001 REGISTERED



DNV

MGMT.SYS.

RvA C:024

MGMT.SYS.

RvA C:425

DNV Certification B.V., The Netherlands

# VAPCO

**VETERINARY & AGRICULTURAL  
PRODUCTS MANUFACTURING CO. LTD.**



## **WARNING:**

If symptoms persist after using this preparation for 2-3 days, consult veterinarian.

## **WITHDRAWAL PERIOD:**

### **Meat consumption:**

Cattle: 1 day.

Sheep: 2 days.

Goat: 3 days.

Poultry: 5 days

## **STORAGE:**

Store in dry place below 30°C.

**PACKING:** (100, 250, 500, 1000) ml.

**FOR VET USE ONLY**