



Phoxivap – 50%

Emulsifiable Concentrate

COMPOSITION: Each ml contains:

Phoxime

500 mg

PROPERTIES:

Phoxime is an inhibitor of the enzyme cholinesterase at the synapses of the nerve ganglion. Inhibition of the enzyme is irreversible under physiological conditions.

INDICATIONS:

Treatment of myiasis, scabies and infestations by ticks, lice and keds in cattle, sheep, goats, horses.

DOSAGE:

By **bathing, spraying or rubbing with a sponge**, in a single application after dilution of the product according to the recommendations below:

	Sheep- Goat	Cattle	Equine
scabies			
Dose	1 Lt in 2000 Lt water. Old scabies: 1 Lt in 1000 Lt water.	1 Lt in 1000 Lt water. Old scabies 1 Lt in 500 Lt water.	1 ml in 1 Lt water. Sponge application.
Keds, Lice, myiasis and Ticks.			
Dose	1 Lt in 2000 Lt water.	1 Lt in 1000 Lt water.	1 ml in 1 Lt water Sponge application.

Replenish: when the volume of the bath has decreased by 20%, refill with a solution of 250 ml of pure product in 125 liters water. Refill the bath the first 3 days and renew the complete bath on the 5th day.

WITHDRAWAL PERIOD:

Cattle:

- Meat and offal: 40 days.
- Milk: do not use in females producing milk for human consumption, lactating or in dry period.

Sheep, goats and horses:

- Meat and offal: 28 days.
- Milk: do not use in females producing milk for human consumption, lactating or in dry period.

ISO 9001 & 14001 REGISTERED



DNV Certification B.V., The Netherlands



MGMT. SYS.
RvA C024



MGMT. SYS.
RvA C425

VAPCO

**VETERINARY & AGRICULTURAL
PRODUCTS MANUFACTURING CO. LTD.**



PRECAUTIONS:

- Do not use the drug 10 days before and 10 days after administration of other cholinesterase inhibitors, phenothiazines or muscle relaxants. Also avoid simultaneous use with general anesthetics.

STORAGE: Keep in dry, dark place below 30°C.

PACKING: (100, 250, 500, 1000) ml.

FOR VET USE ONLY