



# TYLOSIN-50%

Water soluble Powder

### COMPOSITION: Each gm contains:

Tylosin (As Tylosin Tartrate) 500 mg

### INDICATIONS:

For the treatment of chickens, turkeys, and calves suffering from infectious diseases caused by pathogens sensitive to tylosin (chicken and turkey mycoplasmosis, infectious synovitis, borreliosis and necrotic enteritis, turkey infectious sinusitis, calf enteritis and bronchopneumonia).

### DOSAGE:

#### ■ Treatment:

For chickens and Turkey, the drug should be administered orally dissolved in drinking water, for calves in milk or milk replacer. The intended amount of the product, it should first be dissolved in a small amount of water (by pouring water into the powder, and not vice versa), then this solution should be mixed with the required amount of water.

#### ■ Treatment:

- Chickens and turkeys: 1 g of powder should be mixed with 1 liter of drinking water and given 3-5 days.
- Calf: 2 g of powder per animal, 2 times a day, for 7-14 days.

#### ■ Prevention:

For preventive watering of chickens and turkeys, 1 g of powder should be mixed with 1 liter of drinking water according to the following scheme:

Bird species and category	Age of treated birds	Duration of treatment (days)
Broiler chickens	Day 1	3
	Week 3	1
Laying hens	Day 1	3
	Week 3	1
	Week 9	2
	Week 20	2
Breeding flock chickens	Day 1	5
	Week 4	2
	Week 9	2
	Week 16	2
	Week 20	2
	Week 24	2
Broiler turkeys	Day 1	5
	Week 3	1

**WITHDRAWAL PERIOD:****For meat consumption:**

- **Calf:** 14 days.
  - **Chicken :**1 day.
  - **Turkey:** 5 days.
- Do not use in laying birds producing eggs for human consumption.

**CONTRAINDICATION:** Do not use if animals are being vaccinated with live bacterial vaccines at the same time.

**REMARK:**

- The medicated solution should be changed every 24 hours.
- The Product should be dissolved in milk or milk replacer immediately before use.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** (100, 200, 250, 500, 1000) GM.

**FOR VET USE ONLY**