ISO 9001 & 14001 REGISTERED
MGMT.SYS. RvA C024 MGMT.SYS. RvA C425
DNV Certification B.V., The Netherlands

VAPCO
VETERINARY & AGRICULTURAL PRODUCTS MANUFACTURING CO. LTD.

Quality Mark

FIPROVAP 0.25%

Spray

COMPOSITION: Each ml contains:

Fipronil	2.5 mg

INDICATIONS:

Treatment and prevention of infestation by fleas and ticks in dogs and cats. Fipronil exhibits insecticidal and acaricidal activity against fleas (Ctenocephalides spp), ticks (Rhipicephalus spp., Ixodes spp.) and lice (Trichodectes spp. and Felicola spp.), Neotrombicula autumnalis, Sarcoptes spp. and Cheyletiella spp.

DOSAGE AND ADMINISTRATION:

- Spray the entire body of the animal.
- Apply against the lay of the hair and make sure that the entire coat of the animals is wetted.
- Allow to dry naturally. Do not towel dry.

Dosage:

- In order to wet the coat down to the skin, depending on the length of the hair, apply 3 to 6 ml per kg body weight (7.5 to 15 mg of active ingredient per kg body weight).
- The product is active for 4 weeks (3 to 5) against Ticks and 2 months against fleas. (1 to 3 depending on the parasite challenge in the animals environment) in dogs. In cats, the product is active for 6 weeks against fleas.
- Dogs and cats: 6 – 12 spraying / kg.

WARNINGS:

- Avoid Spraying in animals' eyes & directly onto areas of broken skin
- Do not use in rabbits as adverse reactions and even death could occurred
- Wear gloves, especially during treatment animals, and do not smoke, drink or eat during application.
- Keep treated animals away from fires and other sources of heat for at least 30 minutes following spraying and until the fur is totally dry.

STORAGE:

- Keep in a dry, dark place below 30 ºC.

For vet use only