PROFEVAP

(Injectable Solution)
For Cattle and Horses

COMPOSITION: Each 1 ml contains:

Ketorprofen	**100 mg**

INDICATION:

Horse

- The alleviation of inflammation and pain associated with musculoskeletal Disorders;
- The alleviation of visceral pain associated with colic.

Cattle

- The supportive treatment of parturient paresis associated with calving;
- reducing the pyrexia and distress associated with bacterial respiratory disease when used in conjunction with antimicrobial therapy as appropriate;
- improving the recovery rate in acute clinical mastitis, including acute endotoxin mastitis, caused by Gram-negative micro-organisms, in conjunction with antimicrobial therapy;
- reducing oedema of the udder associated with calving.
- Reducing pain associated with lameness.

DOSAGE AND ADMINISTRATION:

Horse:
Intravenous administration.
For use in musculo-skeletal conditions: 1 ml of product per 45 kg body weight, administered by intravenous injection once daily for up to 3 to 5 days.
For use in equine colic: given by intravenous injection for immediate effect. A second injection may be given if colic recurs.

Cattle: Intravenous or intramuscular administration. 1 ml of product per 33 kg body weight, administered by intravenous or deep intramuscular injection once daily for up to 3 days.

CONTRAINDICATION:

- Do not use in animals with known hypersensitivity to the ketoprofen.
- Do not administer other non-steroidal anti-inflammatory drugs (NSAIDs) concurrently or within 24 hours of each other, corticosteroids, diuretics and anticoagulants.
- Do not use in animals suffering from cardiac, hepatic or renal disease, where there is the possibility of gastro-intestinal ulceration or bleeding, or where there is evidence of blood dyscrasia.

WARNINGS:

- The use of ketoprofen is not recommended in foals under the age of 15 days.
- Use in any animal less than 6 weeks of age or in aged animals may involve additional risk. If such use cannot be avoided animals may require a reduced dosage and careful management.
- Should not be administered to pregnant horses.
- Avoid use in any dehydrated, hypovolaemic or hypotensive animals as there is a potential risk of increased renal toxicity.
- Avoid intra-arterial injection.
- Do not exceed the stated dose or duration of treatment.

ADVERSE REACTIONS:

- Mild inflammatory reactions at the injection site.

INTERACTION WITH OTHER MEDICATION:

- Do not administer with other non-steroidal anti-inflammatory drugs (NSAIDs) concurrently or within 24 hours of each other, corticosteroids, diuretics or anticoagulants.
- Concurrent administration with nephrotoxic drugs should be avoided.

WITHDRAWAL PERIOD:

Meat:
Cattle:
Meat and offal: I.V. injection 1 day, I.M. injection 2 days.
Milk: 0 hours.

STORAGE:

Store at a temperature below 30° C.

Keep out of reach and sight of children.

For VET. Use Only