Tilmifen

(Injectable solution)

Composition : Each ml contains :

Tilmicosin (as tilmicosin phosphate)	300 mg
Ketoprofen	100 mg

Indications :

For the treatment and prevention of infectious diseases in cattle caused by organisms sensitive to tilmicosin , mainly respiratory infectious processes , keratoconjunctivitis and foot root , ketoprofen relieves pain ,fever and inflammation associated with any infectious process. It has a high affinity for plasma proteins, which facilitates its passage through the exudates and its accumulation at sites of inflammation

Ketoprofen helps tilmicosin , which has excellent penetration into inaccessible sites, to be used with total effectiveness in the tissues, making the combination of these two actives a therapeutic tool of choice for the treatment of lung , eye, and podal infections

Dosage and administration:

The recommended dose for cattle is 1 ml / 30 kg body weight, one to three applications, 72 hours apart, regarding veterinarian recommendation

Apply only subcutaneously,

Warning and contraindications:

- Do not use in goats
- Do not apply intravenously because it can be fatal
- Do not apply to hypersensitive animal to any component of the product
- Do not use in calves under 1 month or animals which are fed only with milk
-

- The product should not be applied with corticosteroids because it can affect the gastrointestinal mucosa
- The product should not be apply in the last third of gestation because ketoprofen may modify the cardiovascular system of infants
- The product should not be used in animals with kidney or liver diseases
- Do not apply to dehydrated animals , hypotensive , thrombocytopenic or with ulcers or other gastrointestinal disorders
- Should not be used in pregnant cows.

Precautions:

Restrain animals prior to the application. Do not inject more than 20 ml per each injection site, do not mix in the same syringe, tube or container, with any foreign substance to the product (other antibiotics, anti- inflammatories, anesthetics, etc…)

Take precautions to avoid accidental contact or self-injection. Use only subcutaneously, intramuscular injection can cause reactions, wash hands with soap and water after using the product, keep out of reach of children. This product should be applied under the direction of a veterinarian.

Overdose :

In case of overdose, the following symptoms can occur in animals, restlessness, vomiting, and tachycardia and decreased of the cardiac contractility, seizures and death

There is no specific antidote available, apply symptomatic treatment, the administration of intravenous calcium salts compensates the tachycardia

In case of accidental injection or ingestion in human, consult a physician or a center of human toxicology, apply symptomatic treatment, there is no specific antidote

Withdrawal period: discontinue the treatment at least 28 days prior to slaughter, should not be used in lactating dairy cows

Storage conditions:

Keep in dry dark place below 30 c

Packing : (50,100) ml

For vet. use only