VAPCOCOX-FORTE

Oral Solution

COMPOSITION: Each ml contains:

Diclazuril	25 mg

DESCRIPTION:

Diclazuril belongs to the group of benzeneacetonitriles and has anti-coccidial activity against Eimeria species. Depending on the coccidia species, diclazuril has an effect on either the sexual or the asexual stages of the development cycle of the parasite. Diclazuril is poorly absorbed from the intestinal canal, therefore it has mainly effect on coccidia present in the intestines.

INDICATIONS:

Vapcocox-forte is effective in the prevention and treatment of coccidial infections in lambs caused by Eimeria crandalis or Emiria ovinoidalis and in calves caused by Eimeria bovis or Eimeria zuernii. In cases where there is uncertainty whether symptoms are caused by coccidiosis, faeces examination is recommended. Vapcocox-forte oral is effective in the prevention and treatment of coccidial infections in poultry caused by Eimeria tenella, Eimeria necatrix, Eimeria acervulina, Eimeria brunetti, E.mitis (mivati) and E.maxima.

Because diclazuril is effective against E.maxima later in its life cycle, subclinical intestinal lesions maybe present for a short time after infection. Diclazuril was shown in studies to reduce lesions scores and improve performance and health of birds challenged with E.maxima.

DOSAGE:

For oral administration via feed or drinking water. It is advised to treat all animals of the herd at the same time.

Calves, lambs:

Treatment: 1 ml per 25 kg bodyweight in one single treatment.
Prevention: 1 ml per 25 kg bodyweight at about 4 – 6 weeks of age. Under conditions of high infection pressure, a second treatment may be indicated about 3 weeks after the first dose.

Rabbits:

- 0.002 ml per kg bodyweight. This is equivalent to 0.05 mg diclazuril per kg bodyweight.
- In feed: give 4 ml per 100 kg complete feed.

Poultry (broilers):

- Week 1: 30 ml per 1,000 liters of drinking water during 3 days.
- Week 2: 37 ml per 1,000 liters of drinking water during 3 days.
- Week 3: 44 ml per 1,000 liters of drinking water during 3 days.
- Week 4: 51 ml per 1,000 liters of drinking water during 3 days.
- Week 5: 58 ml per 1,000 liters of drinking water during 3 days.
- Week 6: 67 ml per 1,000 liters of drinking water during 3 days.

This is equivalent to 0, 3 mg of diclazuril per kg bodyweight per day.

WITHDRAWAL PERIOD:
Meat: Calves, lambs: 0 days
Poultry: 0 days
Rabbits: 2 days

REMARKS:

- To prevent resistance do not use for a long period, try to rotate with other anti-coccidials treatments.
- Mixed with feed, the product should be used immediately.
- Medicated drinking water should be used within 24 hours.

CONTRA-INDICATIONS:
Do not use in layers producing eggs for human consumption.

STORAGE: Keep in dry, dark place below 30 ºC.

For Vet. Use only.